



- Concentration: 0.6% (6 mg/mL)
- Available sizes: Pint or gallon
- Common prescription dosage:
- Day 1: Two pints of Meloxicam per five gallons of stock solution
- Days 2–7: One pint of Meloxicam per five gallons of stock solution
- Miscibility: Amoxicillin Red, tilmicosin, neomycin, sulfadiazine/ trimethoprim, chlortetracycline, lincomycin, gentamicin, Iversol™
- Do not mix with: Acids, tylvalosin, tiamulin
- Recommended withdrawal period: 17 days\*

ALSO AVAILABLE AS AN ORAL DRENCH PRODUCT.



Have questions? Contact us at:

877-931-8707 • Pharmacy@VetPharmSol.com VeterinaryPharmaceuticalSolutions.com

## ADMINISTERING COMPOUNDED SOLUBLE MELOXICAM



Scan the QR code for an instructional

demonstration.



Meloxicam is miscible with a large variety of other drugs in the same stock solution, including Amoxicillin Red, Iversol,™ tilmicosin, neomycin, sulfadiazine/trimethoprim, chlortetracycline, lincomycin and gentamicin. Mix Meloxicam with other drugs only on veterinarian's prescription.


## **INSTRUCTIONS**



- Use cold water to mix
- Mix precisely to the ratio directed on the prescription 2
- Do not exceed total volume called for on prescription 3
- 4 Add Meloxicam, then water, then mix well with stirring spoon or tool
- 5 Follow duration of treatment per prescription instructions
- 6 Mix Meloxicam with other drugs only on veterinarian's prescription
- 7 Discard old solution and mix new solution daily

## **Have questions? Contact us at:**

877-931-8707 • Pharmacy@VetPharmSol.com VeterinaryPharmaceuticalSolutions.com



Disclaimer: These statements have not been evaluated by the Food and Drug Administration.



https://www.ema.europa.eu/en/medicines/veterinary/EPAR/metacam

https://www.drugs.com/vet/metacam-for-swine-can.html

A study was conducted on growing-age pigs, where a 5x average daily dose was administered for seven days through
a medicator and was fully consumed/disappeared.